

Lit. J. contient soixante et quatre feuilles de toutes sortes de racines, fleurs et

Inhalt 72 Bl. 109 abb.

i

CEPA ASCALONICA
Mattsioli,
filiis & sterilis
Germ. Ascheum.



BVLBVS AGRESTIS Tragi, ac
SILVESTRIS Dodonaeo.

Germ. feldt zwiwe oder Arden zwiwe.
Belg. Wilden Aijnin.



BVLBVS SILVESTRIS
ALTER, flore candido.

1657. 5. Jul. 1602. P. 272. 70.

15. Jul. 1539

2

ΣΧΟΙΝΟΠΡΑΣΟΝ, Dodonæo.
PORRVM IVNCEVM
quibusdam SECTIVVM

Gall. Potit pourreau.
Germ. Signitlaure.
belg. Bijn loorh.

2



Scratur inter olera.
Floret Maio tumiq; mensibus, altero quam satum
aut transplantatum est anno.



3.

ALLIVM VRSINVM Gall. Ail Sauv. forte SCORODOPRASVM Diosc. lib. 2. cap. 145.
 Germ. Wacht Knoblauch Belg. Daelcoock.

ὄφιοσκορόδον
 ALLIVM ANGVINVM
 seu SILVESTRE
 CANINVM PRIMVM Tragi.

Gall. Auy saulvaige.
 Germ. Wilder Knoblauch oder
 feldknoblauch.
 Belg. Wilt Coock of Cravoloo.



Invasitur alicubi segetis. in partem agri
 colis, reperiturq; ad sepes et in pratis
 siccioribus.

Floret semenq; fert mense junio
 julioq;.

Umbris gaudet & sponte nascitur in frator.
 marginibus.
 Floret mense Aprili.

26. 25. 1546.

26. 15. fol. 1547. Dodon. 4. 3. 13.

76





ΣΚΟΡΟΔΟΝ
ALLIVM

Ital. Aglio,
Gall. ail ou auy.
Germ. Knoblauch,
Belg. Coorke.

Diosc. lib. 2. cap. 145.
Theophr. de nat. plant. lib. 7. cap. 4.
Galen. lib. 2. de alim. facult.



Scritur nucleat in divisum in hortis.
Floret

Lib. 15 fol. 154r.



ORCHIS
Flore purp. maculosa
minor.

+

ORCHIS
Flore purpureo,
sepalibus maris
infrimis
ORCHIOS
secundum genus
Dodonæo.

Orchis alba
filio suo maculato.

ORCHIOS
SPECIES
in microscopia



Ed. 15. f. c. 1559.

Ed. 15. fol. 1559. et parat. de Piquet.



SATYRIUM
PALMA
flore pur
Serapias violacea.

BASILICVM seu
CHRISTI
pureo.

PALMA CHRISTI
flore purpu
rascente.

ORCHIS SERAPIAS FEMINA seu
TRIORCHIS ALTERA. *Dolongo.*
flore Crabronem referente.

Serapias belliniana

PALMA CHRISTI
flore
candido.



Hec, ut & superiora, locis siluosis & in pratis
frequentia nobis sunt.
Florent mense Maio tumq[ue].

de reli. specibus videb[itur] d[omi]n[us] lib. 15. fol. 1546.

81
vide doc. 2. 2. 31.

Voss de vinctis hb. 15 Jul 1599. in Reginal.

de omnibus specie orchi Dodon. 2. 2. 30.

ORCHIS muscarum flore raro imitante.



ORCHIS can. flore Aido.



TRAGORCHIS Dodoneo:



Belg. Bogruelstend of Gasfrullous.

+ Phycos minor triphylla.




TESTICVLVS Dodo ODORATVS 1120.

Belg. Cluyt vlandelouyt.



Hic pro ceteris locis aprius gaudet. Floret mensis Augusto.



Locus umbrasis & pratensibus gaudet. Floret Maio Iunioq.

Lepidula odvatis. Dod. 2. 2. 30.

b orchi serapias primus dod. 2. 2. 30. a Lobel fol 214

KOAXIKON: COLCHICVM, seu
BVLBVS AGRESTIS

Ital. Colchico.

Gall. Mont an'gisy ou Colchicu.

Germ. Zwitterloß und Wild Safran oder Herbstbeem.

Belg. Wilder offe Heroungstige Edelweiss.

Diosc. lib. 4. cap. 72.
Galen. lib. 6. simp. medic.



Nascitur in pratis vlgiosis, circa Viliuordiam
Brabantie oppidum, frequens.

Florescunt paulo ante autumnale æquinactium:
sed folia proximo post vere prodeunt, semine
mense Junio maturescente.





COLCHICVM *florens*.



15. fol. 1371







Sisymbrium dolonei hb. 15. fol. 1680.



J. G. R. 1615. p. 1615. 1615.

OPNITHOGALON
ORNITHOGALVM

Ital. Cotte di gallina.
Gall. Cquirus.
Germ. Felgen veldmuff.
Belg. vilt velt angwijn.



Nascitur inter segetes, maxime potissimum in
sabuloso solo.
Folia produnt Martio Aprilique: flores vero Maio:
post tamen autem, herba ipsa disparet.



FVNGI SPONGIOSI SPECIES

Flandris Bult.

Cuius carne exsiccata incensa, et in furnum resoluta
apes suffocantur.



Nascitur pingui ac putri solo, in vetustis demolitorum aedificiorum arceis.

Intra .xx. dierum spacium in hanc aut etiam maiorem
magnitudinem excrevit, posteaq; euellit, et in aere sus-
penditur, ac in tepidum furnum aliquoties conijcitur, ut
in futurum usum arefiat.

MYKHTEΣ
FVNGI

Ital. Fungli.
Gall. Lofirons, Champignons seu
Mourrons.
Germ. Spitzling und Styrwimm.
Belg. Camoungier, off Kaddenboot

19



Diosc. lib. 4. cap. 71.
Galen. lib. 8. simp. med.

Porri Si omnes noxy sunt aut saltem
inutiles et qui minima apti.

Siccis & nemorosis tractus anant, & nascuntur ubiq.



ALFALFA
ASTRODELAS
AZOZELAS

del. ...
del. ...
del. ...
del. ...

del. ...
del. ...
del. ...

del. ...
del. ...

del. ...
del. ...
del. ...



ΑΣΦΟΔΕΛΟΣ
ASPHODELVS
ALBVCVS seu HASTVLA REGIA

Ital. Amphodello.
Gall. Asphodelus.
Germ. Goldwurtz.
Belg. Asphellu.

Diosc. lib. 2. cap. 161.
Galen. lib. 6. simp. medic.
Plin. lib. 21. cap. 17.



Montes siliosos amat. seriturq;
in hortis.

Radix eius Napis modicis similis est,
ut ait Plinius: neq; alia numerosior,
LXXX sepe simul acervatis bulbis.



ASPHODELI CAVLIS

Quem Theophrastus, & fere Graeci principesq; Pytha-
goras ANOEPKON, Latini vero ALBVCM,
ut Plinius ait, appellaverunt.

Floret mense Maio: seminaq; in tumo maturantur.

ASPHO DELVS
ex candido purpurascens.



ASPHODELVS FEMINA Galemi in libr. de alim. facult. Dodoneo.
 Quibusdam nunc BVLBVS VOMITORIVS
 ORNITHOGALON MAIUS.



Asphodelus ut videtur bulbosus tab. 15. fol. 1589.

LILIAGO Cordi, lib. histor. plant. cap. 105.
ASPHODELLI, vt apparet, SPECIES.



Nascitur editis apricisque
parua arbusta.

locis, inota rara &

Lib. 16. fol. 105





Liliosyde de las Phoenicias Pl. 14 fol. 150

Liban. fol. 127

ASPHODELVS LVTEVS *Dalony.*
folijs non deciduis.

27



28 15 fe 1500.

28.



27

101
112



131





ΧΑΜΑΙΒΑΛΑΝΟΣ
 CHAMAEBALANVS seu GLANS TERRAE
 Dodonæ & alij recentioribus.

Germ. Erdnuss, Erdpfleger, und Erdamandol.
 Belg. Aerdnoten, oft Mispel, met sterck.



Sponte proveniunt in arvis & circa sepes in plerisque Hollandiæ &
 quibusdam Brabantie Flandriæ, Oceano vicinis tractibus.
 Florent mense Junio, semina paulo post perficiunt.
 Radices vero à nonnullis mense Maio in cibum effodiuntur.



Cette herbe a esté vendue & achetée au Jardin Royal
 de Paris l'année 1664. comme de y faire voir par le
 rapport de Monsieur de la Roche, & de Monsieur de la
 Roche, & de Monsieur de la Roche, & de Monsieur de la Roche.
 Et la nomme au lieu de Passiflora.



ARI SEMEN
tam viride quam maturum.



Aron
ARON MACVLOSVM

Belg. Gysperwolden ralfs wort.

Lib. 2. fol. 779. Vide Tragum, qui plures Suius enumerat species.



Nascuntur omnes frigidis atq. umbratis locis.
 Folia mense Martio Apriliq. prodeunt: que in tumis
 marcescunt, cauliculum nudum relinquentes, in quo
 semina mense Augusto maturescunt.



Arisarum. cum flore Myconi Angustifolium

Tab. 16. fol. 1600.

DRACUN CVLI MINORIS FLOS



Tab. 16. f. 1402.

ΔΡΑΚΟΝΤΙΑ ΜΙΚΡΑ
DRACUNCVLVS seu SERPENTARIA MINOR Matthioli &
MAIOR Dodonei.

Ital. Draguntea minore.
Gall. Serpente Serpentaire.
Germ. Klegg natterwurtz.
Belg. Epwortelen vnde Drackfey
Wortelen.

Diosc. lib. 2. cap. 161.
Theophr. de nat. plant. lib. 7. cap. 11.



Nascitur alicubi sponte in sepibus: at nobis seritur
in hortis.

Floret mense Julio, fructumq;
perficit Augusto.

hb. 15. Jul. 1602.



Cyclamen







Vide fol. 1605 v. 1606. 26. 14.

CYCLAMINVS
folia Gedeo. racco.



KYKLAMINOS
CYCLAMINVS
RAPVM. TVBER seu VMBILICVS TERRAE
Ital. Cyclamino o Lari porcino.
Gall. Cyclamin ou Pain de porreau.
Germ. Stuckapffel und St. Wundbrut.
Belg. Stuckapfel oder St. Wundbrut.

Diosc. lib. 2. cap. 158.
Galen. lib. 7. simp. medic.

Nascitur in umbrosis, sub arboribus, dumetis ac senticetis silvis.
Mense Augusto, ubi folia perierint, florēt ad finem usque Septembris

ΚΟΤΥΛΗΔΩΝ : COTYLEDON
UMBILICVS VENERIS seu ACETABVLVM

Ital. Copercinale.
Hisp. Vazillos.
Germ. Nabelkraut.
Belg. Navelkrant.

Diosc. lib. 4. cap. 78.
Galen. lib. 7. simp. medic.



Saxosis & maritimis locis
gaudet.

Floret mense junio lulioq.

46

46



119

ΙΡΙΣ
IRIS

Italice, Iris, Giglio azzurro, sive Giglio celeste.
Gallis, Blayvil sive flamor.
Germanis, Blau gilyon sive Himmelstern.
Belgis, Blau colin sive Ciste van onse zov.

Diosc. lib. 1. cap. 1.
Plin. lib. 21. cap. 7.
Theophr. lib. 9. cap. 7. ubi
CONSECATRIX, iuxta Theo-
dori versionem appellatur.
Mattbiolus lib. cap.
Val. Cordus. lib. 2. cap. 39.

Nascitur plurima & optima in Alyria, etiam
sponte: nobis autem sata tantum.
Floret post medium veris.
Semen apud nos non perficit, quare nescio an
Mattbiolus recte eam cum semine de pingat.



47

47



Iris domestica L. 15. fol. 100.

Lucce. iris nostras vulgaris fol. 86.

120








51





53



7

126

54





56



57.



23. 130



U



11

8

132



61

61



139



Iris tuberosa


14

136



65

IRIS SILVESTRA TEMPERATA
Linn. Sp. Pl. 1207
Linn. Bot. Linn. Soc. 1753
Linn. Bot. Linn. Soc. 1753



IRIS SILVESTRA TEMPERATA
Linn. Sp. Pl. 1207
Linn. Bot. Linn. Soc. 1753
Linn. Bot. Linn. Soc. 1753

IRIS SILVESTRIS TENNIFOLIA

Valerij Cordi, ut videtur.
Traagus pag. 702 exhibet nomine
Gladioli caerulei minoris



Nascitur sponte circa Tigurum in Heluetia, ut Gesnerus
ait: caule oblongo, tribus aut quatuor flosculis cerni-
leis supra se invicem seminiferis.
Præter item secundum Rhenum iuxta Opheim et Wormaciam.

iris silvestris altera. matte. lib. il. fol. 150.



Floret quando Lutea.



IRIS LVTEA
pro ACORO à pharmacopis
nostratib. perperam hactenus habita

Flandris: Cisse sine
geseu lisse.

Floret mense Maio, & semen
maturat mense Augusto.

Val. Cordus lib. plantar.
2. cap. 44.

Nascitur passim apud nos, palustribus
& vligmosis locis: sed inutilis plane
atq; modora.



Tab. 1. 139

Bot. Soc. No. 91.

CHAMÆIRIS LVTEA



Calamus aromaticus officinarum.
Acorum verum Manardo



Floret serius priori



Iris vulgaris candida floribus multatis (sive irone) L. 1688. p. 100.

Tab. 11. Iron. 97. p. 85. cart. 1

69

69



de cornu nostris - Gal. hist. lib. 10. fol. 63

de acoro dabo. lib. 16. fol. 1618.

141



CHAMAEIRIS Dodonæi lib. cap.
 SILVESTRIS ALTERA Mattioli. lib. cap.

Hec in iconibus Mattioli caule longiori quam oporteat
 expressa est, quum ipse eandem contra in commentarijs
 suis caule brevi esse testetur. Inter folia n. floret,
 ut Similis non immerito dici possit.

Hanc verò Illyricam esse quidam perperam contendunt,
 quum illa a nostrata, non forma aut specie, sed virtus
 tantum differat.

Floret mensibus Maio Junio.

141

ZYPIS
XYRIS & IRIS SILVESTRIS
SPATULA FOETIDA vulgo

Gall. Beapoul plant.
Germ. Wueste Schwarte und Wande
Belg. Wueste Cuystruyt & Stijer

conservant.
sindon este.

71

Diosc. lib. 4. cap. 20.
Galen. lib. 8. simp. medic.



Floret mense Augusto, semina
perficit Septembri.

Seritur nobis plerumque in hortis:
reperitur tamen alieni sponte proveniens,
in locis aquis ac iuxta sepes, et in silvarum
marginibus.

172

IRIS MARITIMA forte. &
RIZOTOMOS Plinij.

72.



Hec ob flacciditatem et Guinis abundantiam
improbatur: & sane tam ex florum quam
radicis et foliorum colore Lacustris apparet.
Floret

Xyris—lobata; Dalech. lib. 14. fol. 172

174

